



## Estimation of Carbon Footprint in Direct Seeded Rice (*Oryza sativa*) under Rainfed Medium Land Situation

Susmita Das<sup>1\*</sup> and Hirak Banerjee<sup>2</sup>

<sup>1</sup>Dept. of Agronomy, Sri Sri University, Cuttack, Odisha (795 004), India

<sup>2</sup>Dept. of Agronomy, Bidhan Chandra Krishi Viswavidyalaya, Nadia, West Bengal (741 252), India



Open Access

### Corresponding Author

Susmita Das

✉: susmitadas282@gmail.com

**Conflict of interests:** The author has declared that no conflict of interest exists.

### How to cite this article?

Das, S., Banerjee, H., 2024. Estimation of Carbon Footprint in Direct Seeded Rice (*Oryza sativa*) under Rainfed Medium Land Situation. *Innovative Farming* 9(1), 01-06. DOI: 10.54083/IF/9.1.2024/01-06.

**Copyright:** © 2024 Das and Banerjee. This is an open access article that permits unrestricted use, distribution and reproduction in any medium after the author(s) and source are credited.

### Abstract

An experiment was carried out for two successive years, 2018 and 2019 during the *kharif* season on direct seeded rice in the experimental farm of the Central Agricultural University, Imphal. The research site was located under the eastern Himalayan region (II). The research was carried out to investigate into the carbon footprint of direct seeded rice under rainfed medium land situation. The study was based on factorial randomized block design (FRBD) comprising of two aspects, sowing techniques and seed rate. Broadcasting and line sowing were the levels under sowing techniques and seed rate has five levels including 80 kg ha<sup>-1</sup>, 90 kg ha<sup>-1</sup>, 100 kg ha<sup>-1</sup>, 110 kg ha<sup>-1</sup> and 120 kg ha<sup>-1</sup>, respectively. The overall estimated greenhouse gas emission in line sowing method (11.02% was at par with broadcasting method (11.06%) with 120 kg ha<sup>-1</sup> of seed rate followed in both the methods. Line sowing with 100 kg ha<sup>-1</sup> led to the maximum carbon emissions. Further, line sowing with 80 kg ha<sup>-1</sup> was superior in terms of parameters like carbon sustainability index, carbon efficiency and carbon efficiency ratio.

**Keywords:** Carbon footprint, Direct seeded rice, Greenhouse gas, Seed rate, Sowing techniques

### Introduction

Rice (*Oryza sativa* L.) serves as a fundamental staple crop, feeds over half of the world's population. Nearly 12% of the world's arable land is dedicated to rice cultivation (FAO, 2020). Various agricultural operations during rice cultivation has been an essential factor to the greenhouse gas emission in the form of carbon dioxide (CO<sub>2</sub>) from fossil fuels, release of methane gas (CH<sub>4</sub>) due to submergence in traditional tilled-transplanted rice system, cattle rearing and nitrous oxide (N<sub>2</sub>O) from inorganic and natural fertilizer and compost management practices (Pandey and Agrawal, 2014; Tjandra *et al.*, 2016; Ashoka *et al.*, 2017; Yadav *et al.*, 2018). The conventional tilled-transplanted rice production system has been seen to adversely affect the environment and reducing the profitability of rice cultivation. Agriculture alone accounts for 8.8 to 10.2% of total greenhouse gas emission. Globally, rice production systems have purportedly released around 523 million tonnes CO<sub>2</sub>-equivalent of greenhouse

gas year<sup>-1</sup>. Rice is cultivated on 43 million ha in India and studies reported an emission of 96.2 million tonnes of CO<sub>2</sub>-equivalent year<sup>-1</sup>, which was 18.4% to the global greenhouse gas emission from rice fields as per 2016-2017 reports (FAO, 2017). The bed planting method (BP), direct-seeded rice (DSR), zero tillage (ZT) are the other alternatives to traditional tilled-transplanted system of rice which escapes the operations like tillage, puddling, transplanting; hence, would reduce the emission from the inputs required for agricultural produce (Wassmann *et al.*, 2004; Pathak *et al.*, 2011). In India, the direct seeded rice is mostly grown in uplands which cover roughly 4.95 million ha (12%) of total rice area (FAO, 2017).

Till today, majority of the analysis regarding the release of greenhouse gases from direct-seeded rice has been analyzed by Pathak and Wassmann (2007), Ahmad *et al.* (2009), Pathak (2015), Chaudhary *et al.* (2017) and Yadav *et al.* (2017). Keeping this in view, a field based study has been

### Article History

RECEIVED on 21<sup>st</sup> June 2023

RECEIVED in revised form 08<sup>th</sup> January 2024

ACCEPTED in final form 15<sup>th</sup> January 2024



performed to estimate the carbon footprint in direct-seeded rice cultivation in the rainfed area of Imphal, Manipur.

## Materials and Methods

### Characterization of Experimental Site

The study area laid within the eastern Himalayan region (II) and the sub-tropical zone (NEH-4) of Manipur. In the experimental farm of Central Agricultural University, Imphal, a study was conducted for two successive years during the *kharif* season of 2018 and 2019. The coordinates of the study plot was 24.45° N and 93.56° E and elevation of 790 m above mean sea level. The land was moderately leveled with clay textured soil. The pH value of 5.5 of the initial soil analyses documented that the soil was slightly acidic with high organic carbon content of 1.15%. The N availability of the soil was medium (322 kg ha<sup>-1</sup>), the available P<sub>2</sub>O<sub>5</sub> and available K<sub>2</sub>O in the soil was also medium (17.59 kg ha<sup>-1</sup> and 287.17 kg ha<sup>-1</sup>, respectively). The mean temperature during both the years of experiment has recorded 27.63 °C, as a maximum and 18.85 °C, as a minimum. The average yearly rainfall of the site was 1730 mm.

### Experimental Details

The research were laid under factorial randomized block design (FRBD) with two aspects: planting methods with two stage - line sowing and broadcasting, and another aspect was seed rate with five levels, 80 kg ha<sup>-1</sup>, 90 kg ha<sup>-1</sup>, 100 kg ha<sup>-1</sup>, 110 kg ha<sup>-1</sup>, 120 kg ha<sup>-1</sup>. The experiment consisted of 10 treatments replicated thrice. Plot sizes for each treatment are 3 m × 4 m.

The treatment were paired as: S<sub>1</sub>R<sub>1</sub> = Broadcasting + seed rate (80 kg ha<sup>-1</sup>), S<sub>1</sub>R<sub>2</sub> = Broadcasting + seed rate (90 kg ha<sup>-1</sup>), S<sub>1</sub>R<sub>3</sub> = Broadcasting + seed rate (100 kg ha<sup>-1</sup>), S<sub>1</sub>R<sub>4</sub> = Broadcasting + seed rate (110 kg ha<sup>-1</sup>), S<sub>1</sub>R<sub>5</sub> = Broadcasting + seed rate (120 kg ha<sup>-1</sup>), S<sub>2</sub>R<sub>1</sub> = Line sowing + seed rate (80 kg ha<sup>-1</sup>), S<sub>2</sub>R<sub>2</sub> = Line sowing + seed rate (90 kg ha<sup>-1</sup>), S<sub>2</sub>R<sub>3</sub> = Line sowing + seed rate (100 kg ha<sup>-1</sup>), S<sub>2</sub>R<sub>4</sub> = Line sowing + seed rate (110 kg ha<sup>-1</sup>), S<sub>2</sub>R<sub>5</sub> = Line sowing + seed rate (120 kg ha<sup>-1</sup>).

The Tamphaphou paddy variety (CAU R1) having duration of 135-140 days was used in this experiment because of its consumable quality. The field was ploughed thoroughly once by tractor followed by power tiller. It was then leveled and formed to have a submergence condition for rice paddy cultivation. The treatments were arranged according to the design. Potassium in the form of muriate of potash, nitrogen in the form of urea and phosphorus in the form of single super phosphate, at the rate of 30, 40, 60 kg ha<sup>-1</sup>, respectively, were applied in each treatment in split doses. Three separate applications of nitrogen were made; 50% of the nitrogen was applied as basal and the other 50% was splitted into two identical halves, one at the 35 DAS (active tillering stage) and the other at the 65 DAS (bloom initiation stage). The total amount of potash and phosphorus that was advised was applied as basal. The pre-sprouted seeds were broadcasted or line sowed as per the proposed seed rate on second week of June during both the years of experiment.

### Carbon Footprint Estimation

The ecological impact of direct seeded rice cultivation

was estimated by greenhouse gases emission. In the present research, the carbon footprint of direct seeded rice farming from the field to the farm gate was examined in terms of both spatial and yield scales. The sum total of major greenhouse gases like CH<sub>4</sub>, N<sub>2</sub>O and CO<sub>2</sub> emitted throughout the production of a crop when expressed in terms of CO<sub>2</sub> equivalents is referred to as spatial carbon footprint (Pratibha *et al.*, 2016). The equivalent emission coefficients have been illustrated in table 1 (Deng, 1982; Dyer and Desjardins, 2003; Lal, 2004; Tabatabaie *et al.*, 2012; Gathorne-Hardy, 2016; Vetter *et al.*, 2017).

These emissions were estimated as per the standard emission coefficients prescribed by IPCC (2017); where CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O were transformed into equivalence of CO<sub>2</sub> by using conversion factors of 1, 25 and 298, based on volume for CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O, respectively.

The emission of CH<sub>4</sub> gas from partially submerged paddy field and emissions of N<sub>2</sub>O gas from urea fertilizer was represented after some modifications.

Emission of CH<sub>4</sub> = EF × SF × A × D × 10<sup>-6</sup> (Tubiello *et al.*, 2014)  
..... (1)

Table 1: List of carbon dioxide equivalence factors used in direct seeded rice cultivation

Item	Units	Kg CO <sub>2</sub> -e ha <sup>-1</sup>	References
<b>Land Preparation</b>			
Human labor	day	0.86	Deng (1982)
Fuel-diesel	kg	2.68	Deng (1982)
Cultivator	hr	3.70	Dyer and Desjardins (2003)
Disk plough	hr	5.90	Dyer and Desjardins (2003)
Tractor	hr	12.27	Gathorne-Hardy (2016)
Power Tiller	hr	12.27	Gathorne-Hardy (2016)
<b>Chemical Fertilizer</b>			
Nitrogen	kg	1.30	Tabatabaie <i>et al.</i> (2012)
Phosphorus	kg	0.20	Tabatabaie <i>et al.</i> (2012)
Potassium	kg	0.20	Tabatabaie <i>et al.</i> (2012)
<b>Plant Protection Chemicals</b>			
Fungicide	l	3.90	Lal (2004)
Herbicide	l	6.30	Lal (2004)
Insecticide	l	5.10	Lal (2004)
Chemical spray	l	0.70	Lal (2004)
<b>Seeds</b>			
Rice grain	kg	5.65	Vetter <i>et al.</i> (2017)

[NB: hr = hour; kg = kilogram; l = litre]

Where,

EF = Combined methane emission factor emitted per season, 10 g m<sup>-2</sup>year<sup>-1</sup> for India (Conrad *et al.*, 1996; Parashar *et al.*, 1996).

SF = 0.8 for without organic amendment and flood prone rainfed condition (Houghton *et al.*, 1996).

A = Harvested rice paddy area (ha year<sup>-1</sup>).

D = Duration of cultivation (in days).

N<sub>2</sub>O emissions = N × EF<sub>1</sub> × 44/28 ..... (2)

Where,

N<sub>2</sub>O emissions = N<sub>2</sub>O emissions from crop waste additions to managed soils and synthetic nitrogen manure (kg N<sub>2</sub>O year<sup>-1</sup>).

N = Nitrogen consumption through crop residue, manure and fertilisers, etc. (kg N input year<sup>-1</sup>).

EF<sub>1</sub> = For N<sub>2</sub>O emissions from N inputs, the emission factor is 0.01 (kg N<sub>2</sub>O-N kg<sup>-1</sup> N intake)

GWP = (1 × emission of CO<sub>2</sub> + 25 × emission of CH<sub>4</sub> + 298 × emission of N<sub>2</sub>O) ..... (3)

Where,

GWP = Potential for global warming (kg CO<sub>2</sub>-e ha<sup>-1</sup>).

The summation of the potential of global warming values from all the stages gives us an idea of the spatial carbon footprint and yield-based carbon footprint.

Spatial carbon footprint (GWP<sub>s</sub>) was calculated as,

$$GWP_s = \sum_{i=0}^n GWP_i \dots\dots\dots (4)$$

Where,

n = Number of factors that affected the global warming potential values.

i = Starting at the value on the right side of the equation and ending with the value above the summation sign (n), the index assumes values.

Yield-based carbon footprint (GWP<sub>y</sub>) = Spatial carbon footprint Grain yield ..... (5)

By relating grain yield to the system's global warming potential, the yield-based global warming potential (GWP<sub>y</sub>) or greenhouse gas intensity may be estimated, which aids in measuring and identifying the efficiency of any production system.

*Measures of Carbon Input and Output, Carbon Efficiency, Carbon Sustainability Index and Carbon Efficiency Ratio*

The carbon (C) input was estimated as the total carbon emission or the spatial carbon footprint multiplied by the factor 12/44 as suggested by Chaudhary *et al.* (2017). The carbon equivalent of different plant parts like grain, straw plus root biomass of the rice crop when summed together gives the carbon output. The overall carbon present in the whole crop was measured by multiplying the harvest with 40% carbon (assuming that it is present in the plant biomass).

Carbon efficiency and their related parameters used in the

experiment were given by Lal (2004) and Chaudhary *et al.* (2017) as follows:

Carbon input = Overall carbon emission (kg CO<sub>2</sub>-e) from all factors × 12/44 ..... (6)

Carbon output = (Grain yield × Carbon equivalent) + (Straw yield × Carbon equivalent) ..... (7)

Carbon efficiency (CE) = Carbon output / Carbon input ..... (8)

Carbon sustainability index (CSI) = (Carbon output - Carbon input) / Carbon input ..... (9)

Carbon efficiency ratio (CER) = Grain yield in terms of carbon equivalent / Total carbon input ..... (10)

### Statistical Analysis

For testing the importance of the overall variations between the treatments, analysis of variance (ANOVA) was applied wherever appropriate. According to Gomez and Gomez (1984), to determine whether the two treatment mean differences are significant, the critical difference value at P = 0.05 was computed when 'F' value was determined to be noteworthy.

## Results and Discussion

### Carbon Footprint in Direct Seeded Rice Production

The cultivation operations in direct seeded rice with different levels of seed rate under broadcasting and line sowing techniques contributed significantly to the greenhouse gases emissions (represented in Table 2). Sowing of seed at 120 kg ha<sup>-1</sup> under line sowing technique has resulted in maximum CO<sub>2</sub> emission closely followed by broadcasting of same seed rate. The nitrous oxide and methane emission levels were the same for all the treatment combinations. It was seen in the study that varied seed rate has no significant effect on this two greenhouse gases emission. This was due to same rate of chemical fertilizers applied to all the treatments and drainage facility at frequent interval during various growth stages of rice. The level of nitrous oxide emissions was more compared to methane emission. This was due to methane gas being produced by obligate anaerobic bacteria under continuously submerged rice field, the direct seeded rice being sown in well puddled wet-bed under intermittently flooded condition, methane production was low. This study corroborates with Khalil *et al.* (2004) and Wang *et al.* (2017).

However, this alternate drying and wetting of soils leads to larger microbial activity thus enhancing nitrous oxide gas emission (Harrison-Kirk *et al.*, 2013). The treatment with 120 kg ha<sup>-1</sup> in line sowing followed by broadcasting with the same seed rate exhibited 11.02% and 11.06% of total greenhouse gas emission or spatial carbon footprint respectively, which was the highest among all the treatments. This variation in emission was because of more human labor required in line sowing and higher quantity of seeds sowed than the optimum through line sowing than in broadcasting. The carbon foot print study indicated that the CO<sub>2</sub>-e emissions from seeds contributed the maximum followed by human labor and thirdly by fertilizers more specifically in line sowing

Table 2: Greenhouse gases emission as influenced by varied sowing techniques and seed rates in our study of direct seeded rice cultivation (mean data of two experimental years)

Treatment												Total GHG E	CFy (CO <sub>2</sub> -e kg kg <sup>-1</sup> )
ST	SR	Labor	Die-sel	Mach-inery	Fert-ilizer	Pesti-cide	Pesti-cide spray	Seeds	Total CO <sub>2</sub> emission	Total N <sub>2</sub> O emission	Total CH <sub>4</sub> emission		
BC	80	102	56	75	92	10	1.72	452	790	283	9.72	1083	0.21
	90	108	56	75	92	10	1.72	509	852	283	9.72	1145	0.20
	100	119	56	75	92	10	1.72	565	919	283	9.72	1212	0.21
	110	124	56	75	92	10	1.72	622	982	283	9.72	1274	0.23
	120	130	56	75	92	10	1.72	678	1044	283	9.72	1336	0.26
LS	80	108	56	75	92	10	1.72	452	795	283	9.72	1088	0.20
	90	113	56	75	92	10	1.72	509	857	283	9.72	1150	0.20
	100	124	56	75	92	10	1.72	565	925	283	9.72	1218	0.20
	110	130	56	75	92	10	1.72	622	987	283	9.72	1280	0.22
	120	135	56	75	92	10	1.72	678	1049	283	9.72	1342	0.25
Total									9200	2830	97.20	12128	2.19

[NB: ST = Sowing Techniques; SR = Seed rate; BC = Broadcasting; LS = Line sowing; Pesticide = Fungicide + Herbicide + Fungicide; CFs = Spatial carbon footprint; CFy = Yield scaled carbon footprint; Total GHG: Total GHG emission or CFs (CO<sub>2</sub>-e kg ha<sup>-1</sup>)]

method than broadcasting method. The highest carbon footprint in respect of yield (CF<sub>y</sub>) was found in broadcasting method with 120 kg ha<sup>-1</sup> (0.26 kg CO<sub>2</sub>-eq kg<sup>-1</sup> rice), which also followed the similar pattern as of carbon footprint in respect of space (CF<sub>s</sub>). This indicates less efficient rice production system with higher CF<sub>y</sub>. But then again it was much lesser than the annual average of 5.65 kg CO<sub>2</sub>-e kg<sup>-1</sup> rice in Indian agriculture (Vetter *et al.*, 2017). From the experiment, 9200 kg ha<sup>-1</sup> (~ 75%) of total carbon dioxide gas, 2830 CO<sub>2</sub>-e kg ha<sup>-1</sup> (~ 24%) of nitrous oxide gas and 97.20 CO<sub>2</sub>-e kg ha<sup>-1</sup> (0.81%) of methane was released (Table 2).

From the average data of two years, the carbon input and carbon output gave varied result with different seed rate and sowing techniques (data presented in Table 3).

Among all the treatments, the carbon input was lowest in broadcasting technique compared to line sowing technique. The treatment in broadcasting method with 80 kg ha<sup>-1</sup> recorded the lowest carbon input of 295 kg ha<sup>-1</sup> while line sowing method with 100 kg ha<sup>-1</sup> was highest in terms of carbon output (6382 kg ha<sup>-1</sup>). This could be attributed to increased human labor involvement in line sowing technique for the field operations such as sowing, harvesting and threshing. The treatment in line sowing method with 80 kg ha<sup>-1</sup> was superior in terms of carbon sustainability index and carbon efficiency. This higher carbon sustainability index and carbon efficiency in line sowing method was owing to the optimum carbon output (grain yield) although with lesser carbon input as any cropping system becomes sustainable

Table 3: Evaluation of carbon parameters in direct seeded rice cultivation due to varied sowing techniques and seed rate (mean data of two experimental years)

Treatments		Carbon input (kg ha <sup>-1</sup> )	Carbon output (kg ha <sup>-1</sup> )	Carbon efficiency	Carbon efficiency ratio (CER)	Carbon efficiency ratio (CER)
Sowing Techniques	Seed rate (kg ha <sup>-1</sup> )					
Broadcasting	80	295	5,870	20	18.88	7.06
	90	312	6,094	20	18.52	7.23
	100	331	6,176	19	17.68	6.90
	110	348	6,028	17	16.35	6.26
	120	364	5,932	16	15.28	5.67
Line sowing	80	297	6,084	21	19.50	7.52
	90	314	6,162	20	18.65	7.28
	100	332	6,382	19	18.22	7.26
	110	349	6,223	18	16.83	6.53
	120	366	6,118	17	15.72	5.91

with increased efficiency of the inputs. Consistent findings were reported by Lal (2004), Chaudhary *et al.* (2017) and Yadav *et al.* (2018). Thus, it can be concluded from the study that line sowing technique with 100 kg ha<sup>-1</sup> seed rate in direct seeded rice field could be a significant way to cut the potential of global warming of the conventional rice cultivation system.

## Conclusion

According to the evidence from this two-year study on the carbon footprint of direct-seeded rice under rain-fed medium land conditions, it can be concluded that the line sowing technique, particularly at a seed rate of 100 kg ha<sup>-1</sup>, offers a substantial reduction in greenhouse gas emissions compared to traditional broadcasting methods. The line sowing method demonstrated superior performance in context of carbon efficiency, carbon efficiency ratio and carbon sustainability index, highlighting its potential as a sustainable agricultural practice. While both broadcasting and line sowing at a seed rate of 120 kg ha<sup>-1</sup> resulted in the highest greenhouse gas emissions, the overall carbon output was significantly higher with line sowing. The findings emphasize the importance of optimizing seed rates and sowing techniques to improve the sustainability of rice production. Future studies should focus on refining these practices to further mitigate the environmental impact and enhance the efficiency of rice cultivation systems. This study provides a crucial step towards developing sustainable agricultural practices that can contribute to global efforts in reducing the carbon footprint of crop production.

## Acknowledgements

The author sincerely acknowledge the Department of Agronomy, Central Agricultural University, Imphal, Manipur, India, for supporting this research work and giving necessary amenities during the research.

## References

- Ahmad, S., Li, C., Dai, G., Zhan, M., Wang, J., Pan, S., Cao, C., 2009. Greenhouse gas emission from direct seeding paddy field under different rice tillage systems in central China. *Soil and Tillage Research* 106(1), 54-61. DOI: <https://doi.org/10.1016/j.still.2009.09.005>.
- Ashoka, P., Meena, R.S., Gogoi, N., Kumar, S., Yadav, G.S., Layek, J., 2017. Green nanotechnology is a key for eco-friendly agriculture. *Journal of Clean Production* 142(Part 4), 4440-4441. DOI: <https://doi.org/10.1016/j.jclepro.2016.11.117>.
- Chaudhary, V.P., Singh, K.K., Pratibha, G., Bhattacharyya, R., Shamim, M., Srinivas, I., Patel, A., 2017. Energy conservation and greenhouse gas mitigation under different production systems in rice cultivation. *Energy* 130, 307-317. DOI: <https://doi.org/10.1016/j.energy.2017.04.131>.
- Conrad, R., Mitra, A.P., Neue, H.U., Sass, R., 1996. Methane emissions from rice cultivation. In: *Revised IPCC Guidelines for rational greenhouse gas inventories: Reference Manual*, Volume 3. Intergovernmental Panel on Climate Change (IPCC). pp. 4.53-4.71.
- Deng, J.L., 1982. Control problems of grey systems. *Systems & Control Letters* 1(5), 288-294. DOI: [https://doi.org/10.1016/S0167-6911\(82\)80025-X](https://doi.org/10.1016/S0167-6911(82)80025-X).
- Dyer, J.A., Desjardins, R.L., 2003. Simulated farm fieldwork, energy consumption and related greenhouse gas emissions in Canada. *Biosystems Engineering* 85(4), 503-513. DOI: [https://doi.org/10.1016/S1537-5110\(03\)00072-2](https://doi.org/10.1016/S1537-5110(03)00072-2).
- FAO, 2017. *The State of Food and Agriculture: Leveraging Food Systems for Inclusive Rural Transformation*. Food and Agriculture Organization of the United Nations, Rome, Italy. p. 26.
- FAO, 2020. *OECD-FAO Agricultural Outlook 2020-2029*. OECD Publishing, Paris/ Food and Agriculture Organization (FAO) of the United Nations, Rome. p. 330. DOI: <https://doi.org/10.1787/1112c23b-en>.
- Gathorne-Hardy, A., 2016. The sustainability of changes in agricultural technology: The carbon, economic and labor implications of mechanization and synthetic fertilizer use. *Ambio* 45, 885-894. DOI: <https://doi.org/10.1007/s13280-016-0786-5>.
- Gomez, K.A., Gomez, A.A., 1984. *Statistical Procedures for Agriculture Research*. 2<sup>nd</sup> Edition. John Wiley and Sons Publishers, New York. pp. 357-423.
- Harrison-Kirk, T., Beare, M.H., Meenken, E.D., Condon, L.M., 2013. Soil organic matter and texture affect responses to dry/wet cycles: Effects on carbon dioxide and nitrous oxide emissions. *Soil Biology and Biochemistry* 57, 43-55. DOI: <https://doi.org/10.1016/j.soilbio.2012.10.008>.
- Houghton, J.T., Meira Filho, L.G., Callander, B.A., Harris, N., Kattenberg, A., Maskell, K., 1996. *Climate Change 1995: The Science of Climate Change*. Contribution of Working Group I to the Second Assessment Report of the Intergovernmental Panel on Climate Change. Intergovernmental Panel on Climate Change (IPCC), Cambridge University Press, Cambridge, UK. p. 572.
- IPCC, 2017. IPCC Emission Factor Database (EFDB). In: *IPCC TFI Side-event at the COP-23*. Bonn, Germany, Cambridge University Press, Cambridge, UK.
- Khalil, K., Mary, B., Renault, P., 2004. Nitrous oxide production by nitrification and denitrification in soil aggregates as affected by O<sub>2</sub> concentration. *Soil Biology and Biochemistry* 36(4), 687-699. DOI: <https://doi.org/10.1016/j.soilbio.2004.01.004>.
- Lal, R., 2004. Carbon emissions from farm operations. *Environment International* 30(7), 981-990. DOI: <https://doi.org/10.1016/j.envint.2004.03.005>.
- Pandey, D., Agrawal, M., 2014. Carbon footprint estimation in the agriculture sector. In: *Assessment of Carbon Footprint in Different Industrial Sectors*, Volume 1. EcoProduction. Springer, Singapore. pp. 25-47. DOI: [https://doi.org/10.1007/978-981-4560-41-2\\_2](https://doi.org/10.1007/978-981-4560-41-2_2).
- Parashar, D.C., Mitra, A.P., Gupta, P.K., Rai, J., Sharma, R.C., Singh, N., Koul, S., Ray, H.S., Das, S.N., Parida, K.M., Rao, S.B., Kanungo, S.P., Ramasami, T., Nair, B.U., Swamy, M., Singh, G., Gupta, S.K., Singh, A.R.,

- Saikia, B.K., Batua, A.K.S., Pathak, M.G., Iyer, C.S.P., Gopalakrishnan, M., Sane, P.V., Singh, S.N., Banerjee, R., Sethunathan, N., Adhya, T.K., Rao, V.R., Palit, P., Saha, A.K., Purkait, N.N., Chaturvedi, G.S., Sen, S.P., Sen, M., Sarkar, B., Banik, A., Subbaraya, B.H., Lal, S., Venkatramani, S., Lal, G., Chaudhary, A., Sinha, S.K., 1996. Methane budget from paddy fields in India. *Chemosphere* 33(4), 737-757. DOI: [https://doi.org/10.1016/0045-6535\(96\)00223-8](https://doi.org/10.1016/0045-6535(96)00223-8).
- Pathak, H., Wassmann, R., 2007. Introducing greenhouse gas mitigation as a development objective in rice-based agriculture: I. Generation of technical coefficients. *Agricultural Systems* 94(3), 807-825. DOI: <https://doi.org/10.1016/j.agsy.2006.11.015>.
- Pathak, H., Saharawat, Y.S., Gathala, M., Ladha, J.K., 2011. Impact of resource-conserving technologies on productivity and greenhouse gas emission in rice-wheat system. *Greenhouse Gases: Science and Technology* 1(3), 261-277. DOI: <https://doi.org/10.1002/ghg.27>.
- Pathak, H., 2015. Greenhouse gas emission from Indian agriculture: Trends, drivers and mitigation strategies. *Proceedings in Indian National Science Academy* 81(5), 1133-1149. DOI: <https://doi.org/10.16943/ptinsa/2015/v81i5/48333>.
- Pratibha, G., Srinivas, I., Rao, K.V., Shanker, A.K., Raju, B.M.K., Choudhary, D.K., Rao, K.S., Srinivasarao, C., Maheswari, M., 2016. Net global warming potential and greenhouse gas intensity of conventional and conservation agriculture system in rainfed semi-arid tropics of India. *Atmospheric Environment* 145, 239-250. DOI: <https://doi.org/10.1016/j.atmosenv.2016.09.039>.
- Tabatabaie, S.M.H., Rafiee, S., Keyhani, A., 2012. Energy consumption flow and econometric models of two plum cultivars productions in Tehran Province of Iran. *Energy* 44(1), 211-216. DOI: <https://doi.org/10.1016/j.energy.2012.06.036>.
- Tjandra, T.B., Ng, R., Yeo, Z., Song, B., 2016. Framework and methods to quantify carbon footprint based on an office environment in Singapore. *Journal of Cleaner Production* 112(Part 5), 4183-4195. DOI: <https://doi.org/10.1016/j.jclepro.2015.06.067>.
- Tubiello, F.N., Condor-Golec, R.D., Salvatore, M., Piersante, A., Federici, S., Ferrara, A., Rossi, S., Flammini, A., Cardenas, P., Biancalani, R., Jacobs, H., Prasula, P., Prosperi, P., 2014. *Estimating Greenhouse Gas Emissions in Agriculture: A Manual to Address Data Requirements for Developing Countries*. Food and Agriculture Organization of the United Nations, Rome. p. 181.
- Vetter, S.H., Sapkota, T.B., Hillier, J., Stirling, C.M., Macdiarmid, J.I., Aleksandrowicz, L., Green, R., Joy, E.J.M., Dangour, A.D., Smitha, P., 2017. Greenhouse gas emissions from agricultural food production to supply Indian diets: Implications for climate change mitigation. *Agriculture Ecosystems and Environment* 237, 234-241. DOI: <https://doi.org/10.1016/j.agee.2016.12.024>.
- Wang, C., Lai, D.Y.F., Sardans, J., Wang, W., Zeng, C., Penuelas, J., 2017. Factors related with CH<sub>4</sub> and N<sub>2</sub>O emissions from a paddy field: clues for management implications. *PLoS ONE* 12(1), e0169254. DOI: <https://doi.org/10.1371/journal.pone.0169254>.
- Wassmann, R., Neue, H.U., Ladha, J.K., Aulakh, M.S., 2004. Mitigating greenhouse gas emissions from rice-wheat cropping systems in Asia. *Environment, Development and Sustainability* 6, 65-90. DOI: <https://doi.org/10.1023/B:ENVI.0000003630.54494.a7>.
- Yadav, G.S., Lal, R., Meena, R.S., Datta, M., Babu, S., Das, A., Layek, J., Saha, P., 2017. Energy budgeting for designing sustainable and environmentally clean/safer cropping systems for rainfed rice fallow lands in India. *Journal of Cleaner Production* 158, 29-37. DOI: <https://doi.org/10.1016/j.jclepro.2017.04.170>.
- Yadav, G.S., Das, A., Lal, R., Babu, S., Meena, R.S., Saha, P., Singh, R., Datta, M., 2018. Energy budget and carbon footprint in a no-till and mulch based rice-mustard cropping system. *Journal of Cleaner Production* 191, 144-157. DOI: <https://doi.org/10.1016/j.jclepro.2018.04.173>.