

REVIEW ON *CLUPISOMA GARUA* (HAMILTON, 1822), AN INHABITANT SPECIES IN INLAND OPEN WATERS OF INDIA

Dibakar Bhakta* and Sonia

ICAR-Central Inland Fisheries Research Institute, Regional Center, B-12, Hans Society, Harney Road, Vadodara-390 022, Gujarat, INDIA

*Corresponding author's E-mail: dibakarbhakta@gmail.com

KEYWORDS:

Biological information, *Clupisoma garua*, Conservation

ARTICLE INFO

Received on:

27.10.2019

Revised on:

21.02.2020

Accepted on:

28.02.2020

ABSTRACT

Clupisoma garua (Hamilton, 1822) commonly known as 'garuabachcha' under the order Siluriformes and family Schilbeidae, a potamodromous species, demersal habitat and found both fresh as well as brackish water. The species is mainly distributed in Asian countries like Bangladesh, India, Pakistan and Nepal and found mainly large freshwater bodies. The species are categorized as Least Concern (LC) by IUCN but kept as Vulnerable (Vu) category by the CAMP and CAFF reports. It is an omnivorous species and feeds on insects, molluscs, crustaceans and small fishes in wild conditions. The species have both food and ornamental values and the medium-size individual also treated as game fish in India. Through the species is said to be widely distributed but threatened in some localities like southern West Bengal, due to overfishing and anthropogenic activities. In the present review, we have tried to give detail information of 'garuabachcha', which will be useful to know the species, its potential importance and suitable conservation measures may carry out accordingly.

INTRODUCTION

Clupisoma garua (Hamilton, 1822) commonly known as 'garuabachcha' under the order Siluriformes and family Schilbeidae well distributed in Indian rivers and reservoirs and have potential aquaculture importance. The catfish family Schilbeidae native to Africa and Asia comprises of 66 species in 14 genera (Eschmeyer and Fong, 2016). Approximately half (32 species) is known from Asian countries comprising of 5 genera including *Clupisoma*, *Ailia*, *Horabagrus*, *Lalates* and *Pseudeutropius* (Wang *et al.*, 2016). The genus *Clupisoma* are having of five species of which four are reported from Indian region namely *C. garua*, *C. bastari*, *C. naziri* and *C. montana* (Hora, 1937). As per IUCN Red List Status, the species categorized under Least Concern (LC), according to CAMP (1998) and CAFF (2006) report the species declared as Vulnerable (Vu) category due to the reduction of populations in natural habitats. In the neighboring country, Bangladesh the species in recently kept under critically endangered (Hanif *et al.*, 2015; IUCN, 2015) of limited geographical distribution in that region over the decades for the increase of anthropogenic and natural hazards (Siddik *et al.*, 2017).

Clupisoma garua is said to be commercially important freshwater fish species and a potential candidate species for aquaculture system (Saraswat *et al.*, 2014) and its demand is exceeding its supply in the market (Verma and Serajuddin, 2017). The 'garuabachcha' a very important capture fishery

resources and well distributed in Indian rivers and reservoirs and have potential aquaculture importance (Figure 1). Being boneless, it is mostly preferred as food fish (Talwar and Jhingran 1991). The fish is consumed by all groups of consumers but mostly favoured by poor people due to their high nutritional value with protein content of 18.40%, fat 5.2% and water 74.2% (Bhuiyan, 1964; Memon *et al.*, 2010), availability (Galib *et al.*, 2009) and good taste (Siddik *et al.*, 2017; Verma and Serajuddin, 2017). The fish also has an ornamental value hence, promotes livelihood alternative for coastal communities (Gupta *et al.*, 2016). Through the species is said to be widely distributed but threatened in some localities like southern West Bengal, due to overfishing activities (Verma *et al.*, 2014) and a moderate decline of this species from natural water bodies of Bengal also reported by Patra *et al.* (2005) and Mishra *et al.* (2009). This is a migratory species and used to travel for a long distance for feeding as well as to searching for suitable breeding grounds.

Common names: Garua Bachcha, Guarchcha (English).

Bangladesh: Ghaira, Gang ghaira, Ghaura and river catfish (Galib, 2008; Rahman, 1989).

India: Neria (Assam); Garua, Gharwya, Kocho, Puttoshi, Gaurcha, Ghero (West Bengal); Bachawa, Gharuwa (Bihar); Buchua (Uttar Pradesh); Chellee, Buchua (Punjab); Punia-cathua, Gajri (Orissa) (Talwar and Jhingran, 1991).

Finland: Sillimomni (Finnish) (FishBase.org.in)

Distinguishing Characters

Morphological characters are well documented by Jayaram, 1977; Talwar and Jhingran, 1991; Sarma *et al.*, 2009; Das *et al.*, 2012; Gupta and Banerjee, 2016; Hadiuzzaman *et al.*, 2017; Singh *et al.*, 2018.

Fin formula: D.1/6; A 3/28; P. 1/11; V. 1/5; C. 17-20.

Clupisoma garua is herring-shaped fish, tapering very gradually toward both ends and abdominal edge partly keeled between pelvic fins and vent (Jayaram, 1977). The body is elongated and laterally compressed with a silvery body and yellowish-green back. Head is oval, blunt and of moderate size with a golden gloss (Talwar and Jhingran, 1991). Eyes are large with circular adipose eyelids. The mouth is wide and sub-terminal. Teeth villiform on jaws

and palate. Upper jaw is longer. Barbels 4 pairs, 2 pairs of mandibular and one each of maxillary and nasal. Maxillary reaching the middle or to the base of pelvic fin. Mandibular barbels extending to the base of pectoral fins in adults. Nasal barbel not reaching the anterior edge of the eye. Pectoral fin serrated internally and not extending pelvic fin. Dorsal spine is slender. The adipose dorsal fin is absent in adults. Anal fin with 29 to 36 branched rays. Caudal fin deeply forked, lower lobe longer than upper. Dorsal, pectoral and caudal are black edged.

Colour: Body color is silvery grey on the back and lighter on the sides and abdomen; fins are tinted grey in color with dorsal, pectoral and caudal fin are black edged. The species name *garua* probably derived from *gerua* (yellowish orange) color fins.

Maximum length: 60.9 cm of SL (Shrestha, 1990) and 100 cm of TL (Talwar and Jhingran, 1999).



Figure 1: A fully matured *C. garua* collected from a large reservoir resources

Distribution

Global Distribution

Clupisoma garua has a wide distribution, Indus plain and adjacent hilly areas, Pakistan, in the large reservoir of northern India, Bangladesh and Nepal (Talwar and Jhingran, 1991).

Local Distribution

In India, its distributions are to north India including Bihar, West Bengal, Odisha, Madhya Pradesh, Assam (Brahmaputra and Barak drainage).

Habit and Habitat

It is mainly found in lacustrine habitat in normally at larger rivers and reservoirs and also in stagnant impoundments. Generally inhabitant in freshwater body, brackish water

abundance also reported and a bottom dweller species. They are generally moved in shoals in canals, rivers, streams, lakes, reservoirs and swamplands of both freshwater and brackish-water bodies (Verma and Serajuddin, 2017). Potamodromous, migrating within streams, rivers and travels a long distance (more than 100 km) for feeding or for locating suitable breeding grounds in new water bodies to avoid stress conditions of existing habitat (Froese and Pauly, 2013 and Saraswat *et al.*, 2014).

Food and Feeding Habits

It is an Omni-carnivore fish feeding mostly along the bottom and along margins of the river. Being non-selective feeder they subside on anything and everything available in the habitat. As per the classification of Nikolsky (1963), *Clupisoma garua* falls in the category of euryphagous fish. Nath, 1994 reported this fish as highly carnivorous and

predacious in nature. Afsar (1990) studied on food and feeding habits of *Clupisoma garua* and found continuous in its feeding and reported high feeding intensity during September to October. Garuabachcha feeds actively in April to June and September to December (Chondar, 1999). Surface feeding is during the rainy season and bottom feeding during winters (Krishna Murti *et al.*, 1991). In river Brahmaputra, this fish is caught by using goat liver mixed with fats of Ganges river dolphin, *Platynista gangetica* (Roxburgh, 1801) as bait by Fishers of Bin community, Assam (Ahmed *et al.*, 2018). A comparative food and feeding habits of *C. garua* worked on different workers provided in Table 1.

Table 1: Comparative food and feeding habits of *C. garua* by different researchers

Author	Feeding habits
Khan, 1934	Insects and their larvae, crustaceans and small fishes.
Hora, 1937	Preys on crabs, shrimps, fish, insects etc., and vegetable matter.
Karamchandani, 1957	Carp fry and insects.
Motwani and Karamchandani, 1958	Insects of both aquatic and terrestrial, fish, animal flesh, molluscs, crustacean, algae fragments of higher plants, debris and mucus.
Agarwal and Tyagi, 1969	Omnivorous feeding on small fishes, crustacean and pieces of plants.
Tandon <i>et al.</i> , 1977	Bottom and marginal feeder. Smaller size groups are omnivorous, with an increase in size becomes insectivorous and piscivorous.
Talwar and Jhingran, 1991	Molluscs, insects, small fishes and decaying matter.

Clupisoma garua relies largely on smell, taste and touch and on lateral line sense organ to locate and catch the prey. They wait till animal passes by and then dart out to grasp it (Talwar and Jhingran 1991). The fishes have better developed olfactory organ compared to vision (Ghosh, 2018). There is mixed opinion regarding the feeding behavior of *C. garua*. Several workers reported it is an omnivorous in feeding habits (Tandon *et al.*, 1977; Agarwal and Tyagi, 1969) and few of them also be described the species as carnivorous (Khan, 1934; Molur and Walker, 1998).

Reproduction

Clupisoma garua is bisexual in nature and the male and female individuals could be easily identified once they matured with the secondary sexual characters developed

during spawning season. The pectoral fin spine is found longer in male compared to the female during the breeding season. The species become sexually matured at the end of first year and male matured earlier than that of female (Chondar, 1999). It is reported to move in shoals during breeding period and breeds in shallow water during the month of March-August. The breeding also reported in monsoon floods between May to August at Gangetic Brahmaputra system (Chondar, 1999). Bhuiyan (1964) have reported the suitable spawning temperature of this species is between 15.6-20°C at Bangladesh water bodies. It is an oviparous species, fertilization is external in open water and eggs are remained unguarded by the parents (Das *et al.*, 2012).

Conservation Status

As per IUCN Red List Status, the species is categorized as Least Concern, but it is kept as critically endangered in Bangladesh (IUCN, 2000) and Vulnerable in India (Lakra *et al.*, 2010, CAMP Report, 1998). In several parts of the country, the species population is reported to be in decline from natural water resources. The causes for declining of the species could be over exploitation, loss of habitat, human interference, climate change, pollution, siltation and trade but the primary reason was found to be overfishing (Biswas *et al.*, 2018). Knowledge of certain biological aspects of this commercially important species will be useful to conserve this species in natural habitats and food and feeding behaviors along with reproductive traits will help while aquaculture practices.

Conflict of Interest

There are no conflicts of interest and the authors are responsible for the content and of the writing of the manuscript.

REFERENCES

- Afsar, M.R. 1990. Food and feeding habits of a teleostean fish (Ham.) in the Ganga river system. *J. Freshwater Biology*, 2: 159-167.
- Agarwal, V.P. and A.P. Tyagi. 1969. Food and feeding habits and the alimentary canal of freshwater fishes of Muzaffarnagar. *Agra University Journal of Research (Science)*, 18 (1): 15-28.
- Ahmed, I., S. Borah, B.K. Bhattacharjya, A.T. Landge, A. Kakati, B.J. Saud, B.R. Chetia, M. Kumar and L.K.P.R. Dutta. 2018. An indigenous predatory fish catching technique of lower Brahmaputra valley, Assam. *Journal of Entomology and Zoology studies*, 6(3): 491-494.
- Bhuiyan, A.L. 1964. *Fishes of Dacca, Asiat. Soc. Pakistan*, Pub. 1, No. 13, Dacca, p. 71.

- Biswas, I., T.S. Nagesh, A.M. Sajina and R. Kalidoss. 2018. Morphometric variations and meristic counts of *Clupisoma garua*: an approach for assessing stock structure in the river Ganga. *Journal of Experimental Zoology, India*, 21(2): 813-822.
- CAFF, 2006. Conservation Assessment of Freshwater Fish Diversity for Central India. Proc. workshop, organized by National Bureau of Fish Genetic Resources, 25 Nov, CIAE, Bhopal, MP, India.
- CAMP, 1998. Report of the workshop “Conservation, Assessment and Management Plan for Freshwater fishes of India 1997” organized by Zoo Outreach Organization (ZOO) and National Bureau of Fish Genetic Resources, Lucknow, held at NBFGR, September 1997. pp. 156
- Chondar, S.L. 1999. Biology of Finfish and Shellfish. SCSC Publishers, India. pp.514
- Das, M.K. 2007. Environment and fish health: a holistic assessment of inland fisheries in India. Pp. 137-151. In: Goswami U.C. (ed.) Natural and anthropogenic hazards on fish and fisheries. Narendra Publishing House, New Delhi.
- Das, M.K., M.K. Bandopadhyay, A.P. Sharma, S.K. Paul and S. Bhowmick. 2012. Piscine Diversity of River Brahmani-A Checklist (CIFRI Bulletin no. 175).
- Fricke, R. and W.N. Eschmeyer. 2016. Guide to Fish Collections. left angle bracket <http://researcharchive.calacademy.org/research/ichthyology/catalog/collections.asp> right angle bracket. Electronic version.
- Froese, R. and D. Pauly. 2013. FishBase. World Wide Web electronic publication. <http://www.fishbase.org.in>.
- Galib, S.M., M.A. Samad, A.B.M. Mohsin, F.A. Flowra and M.T. Alam. 2009. Present Status of Fishes in the Chalan Beel- the Largest Beel (Wetland) of Bangladesh. *International Journal of Animal Fish Science*, 2:214-218.
- Galib, S.M. 2008. A Study on Fish Diversity and Fishing Gears of Chalan Beel with Reference to Preservation of Catches, Honors dissertation submitted to the Department of Fisheries, University of Rajshahi, Bangladesh, pp.172.
- Ghosh, S.K. 2018. Histological characterization of the olfactory organ in schilbid catfish, *Clupisoma garua* (Hamilton, 1822). *International Journal of Aquatic Biology*, 6 (5): 281-287.
- Gupta, S. and S. Banerjee. 2016. A note on *Clupisoma garua* (Hamilton, 1822), a freshwater catfish of Indian subcontinent (Teleostei: Siluriformes). *Iranian Journal of Ichthyology*, 3(2): 150-154.
- Gupta, S., S.K. Dubey, R.K. Trivedi, B.K. Chand and S. Banerjee. 2016. Indigenous ornamental freshwater ichthyofauna of the Sundarban Biosphere Reserve, India: status and prospects. *Journal of Threatened taxa*, 8(9): 9144-9154.
- Hadiuzzaman, M., M.F. Ali and M.A.B. Habib. 2017. Etymological history of some freshwater fishes of Bangladesh: A review. *International Journal of Fisheries and Aquatic Research*, 2: 24-30.
- Hanif, M.A., M.A.B. Siddik, M.R. Chaklader, A. Nahar and S. Mahmud. 2015. Fish diversity in the southern coastal waters of Bangladesh: present status, threats and conservation perspectives. *Croatian Journal of Fisheries*, 73(4): 251-271.
- Hora, S.L. 1937. The game fishes of India. III. “Garubachcha or Gaurchcha”, *Clupisoma garua* (Hamilton) and two allied species. *Journal of the Bombay Natural History Society*, 39(4): 659-678.
- IUCN. 2000. Red book of threatened fishes of Bangladesh. *IUCN-The world conservation union xii*. <http://en.bdfish.org/2011/05/garua-bachcha-clupisoma-garua-hamilton-1822/>.
- IUCN. 2015. The IUCN Red List of Threatened Species. Version 2015-4. <http://www.iucnredlist.org/details/166588/0>. Accessed on 2016-01-26.
- Jayaram, K.C. 1977. Aid to the identification of siluroid fishes of India, Burma, Sri Lanka, Pakistan and Bangladesh. 2. Siluridae, Schilbeidae, Pangasidae, Amblycipitidae, Akysidae. Records of the Zoological Survey of India, Miscellaneous Publication, Occasional Paper no. 10, 1-33.
- Karamchandani, S.J. 1957. On the occurrence of associates of carp fry in the fry-collection nets and the destructive role played by predatory fish. *Indian Journal of Fisheries*, 4: 47-61.
- Khan, H. 1934. Habits and habitats of food fishes of the Punjab. *Journal of Bombay Natural History Society*, 37: 655-668.
- Krishna Murti, C.R., K.S. Bilgrami, T.M. Das and R.P. Mathur. 1991. The Ganga: A scientific study.
- Lakra, W.S., U.K. Sarkar, R.S. Kumar, A. Pandey, V.K. Dubey and O.P. Gusain. 2010. Fish diversity, habitat ecology and their conservation and management issues of a tropical River in Ganga basin, India. *The Environmentalist*, 30(4): 306-319.
- Memon, N.N., F.N., F.L. Talpur and M.I. Bhangar. 2010. A comparison of proximate composition and fatty acid

- profile of Indus river fish species. *International Journal of Food Properties*, 13(2): 328-337.
- Mishra, S.S., S.K. Acherjee and S.K. Chakraborty. 2009. Development of tools for assessing conservation categories of siluroid fishes of fresh water and brackish water wetlands of South West Bengal, India. *Environmental Biology of Fishes*, 84(4): 395-407.
- Molur, S. and S. Walker. 1998. Report of the Workshop on "Conservation assessment and management plan for freshwater fishes of India".
- Motwani, M.P. and S.J. Karamchandani. 1958. A note on the food and feeding habits of *Clupisoma garua* (Hamilton) and *Eutropiichthysvacha* (Hamilton) from Allahabad (UP). *Current Science*, 27(2): 55-56.
- Nath, S. 1994. Studies on the bioecology of fishes of Jammu Province (Jammu and Kashmir State) India, Part II: Feeding Ecology. *Recent Advances in Fish Ecology, Limnology and Eco-conservation*, 3: 63.
- Nikolsky, G.V. 1963. "The Ecology of Fishes", Academic Press, London, pp. 1-352.
- Patra, M.K., S.K. Acharjee and S.K. Chakraborty. 2005. Conservation categories of siluroid fishes in North-East Sundarbans, India. *Biodiversity and Conservation*, 14: 1863-1876.
- Rahman, A.K.A. 1989. *Freshwater Fishes of Bangladesh*, 1st edition, Zoological Society of Bangladesh, Department of Zoology, University of Dhaka, Dhaka-1000, p. 185.
- Saraswat, D., W.S. Lakra, P. Nautiyal, M. Goswami, K. Shyamakant and A. Malakar. 2014. Genetic characterization of *Clupisoma garua* (Hamilton, 1822) from six Indian populations using mtDNA cytochrome b gene. *Mitochondrial DNA*, 25(1): 70-77.
- Sarma, D., P.C. Mahanta, D. Sarma and A. Dutta. 2009. Coalmines degraded Ichthyofaunal Diversity of Simsang River, Meghalaya. A report on climate change. Bulletin no-15. Published by Directorate of Coldwater Fisheries Research (Indian Council of Agricultural Research). Bhimtal, Nainital.
- Shrestha, T.K. 1990. Resource ecology of the Himalayan waters. Curriculum Development Centre, Tribhuvan University, Kathmandu, Nepal. 645 p.
- Siddik, M.A.B., M.R. Chaklader, M.A. Hanif, A. Nahar, I. Ilham, A. Cole and R. Fotedar. 2017. Variation in the life-history traits of a Schilbid catfish, *Clupisoma garua* (Hamilton, 1822) in the coastal waters of southern Bangladesh. *Chinese Journal of Oceanology and Limnology*, 35(5): 1189-1196.
- Singh, A.D., S. Abujam and D.N. Das. 2018. *Biodiversity of Fishes in Arunachal Himalaya: Systematics, Classification, and Taxonomic Identification*. Academic Press.
- Talwar, P.K. and A.G. Jhingran. 1991. *Inland Fishes of India and Adjacent Countries*, Vol. II, Oxford & IBH Publishing Co. Pvt. Ltd. New Delhi-Calcutta, pp. 596-597.
- Tandon, K.K., M.S. Johaland and S.K. Dhawan. 1977. Food and feeding habits of some fishes of the Punjab. Punjab Fisheries Bulletin, 1(4): 3-13.
- Verma, J. and M. Serajuddin. 2017. Intra-specific and inter-generic phylogenetic relationships in endangered catfish (*Clupisoma garua* and *Eutropiichthysvacha*) of family schilbeidae. *Journal of Entomology and Zoology Studies*, 5: 198-202.
- Verma, J., A. Kashyap and M. Serajuddin. 2014. Phylogeny based on truss analysis in five populations of freshwater catfish: *Clupisoma garua*. *International Journal of Science and Research*, 3(8): 1414-1418.
- Wang, J., B. Lu, R. Zan, J. Chai, W. Ma, W. Jin, R. Duan, J. Luo, R.W. Murphy, H. Xiao and Z. Chen. 2016. Phylogenetic Relationships of Five Asian Schilbid Genera Including *Clupisoma* (Siluriformes: Schilbeidae). *PloS one*, 11(1): p.e0145675.

How to cite this article?

Bhakta, D., Sonia., 2020. Review on *Clupisoma garua* (Hamilton, 1822), an inhabitant species in inland open waters of India. *Innovative Farming* 5(1): 25-29.