



Short Communication

Article: SAAI/IF/403

Effect of Different Weed Management Practices on the Weed Population Dynamics of Direct Seeded Upland Rice under Tripura Condition

Mandira Chakraborti^{1*}, Buddhadeb Duary² and Mrinmoy Datta^{3,4}

¹Subject Matter Specialist (Agronomy), Krishi Vigyan Kendra (ICAR), West Tripura, Champaknagar, Tripura (799 045), India

²Associate Professor, Dept. of ASEPAN, Palli Siksha Bhavana, Visva-Bharati, Sriniketan, West Bengal (731 236), India

³Former Joint Director, ICAR (RC) for NEH Region, Tripura Centre, Lembucherra, Tripura (799 210), India

⁴Principal, College of Agriculture, Tripura, Lembucherra, West Tripura (799 210), India

*Corresponding email: cmandira1@rediffmail.com

KEYWORDS:

Direct seeded rice,
Economics, Hand
weeding, Pendimethalin,
Weed management

ARTICLE INFO

Received on:

12.07.2020

Revised on:

23.01.2021

Published on:

29.01.2021

ABSTRACT

A field experiment was conducted at Krishi Vigyan Kendra, South Tripura during the *kharif* (wet) season of 2013 and 2014 to study the influence of different weed management practices on weed population dynamics of direct seeded upland rice under Tripura condition. The experiment consisted of twelve treatments laid out in randomized complete block design with three replications. Among the various treatments, ICAR(RC) for NEH Region, Umiam, Meghalaya, pendimethalin at 1.0 kg ha⁻¹ at 2 DAS + bispyribac sodium at 25 g ha⁻¹ at 20 DAS (T₄) recorded the lowest number of all weed in both the years followed by pendimethalin at 1.0 kg ha⁻¹ at 2 DAS + one hand weeding at 30 DAS (T₃). At 60 and 90 DAS, lowest weeds population was recorded by hand weeding thrice at 15, 30 and 45 DAS (T₁₁). The highest net return and benefit cost ratio was observed with the treatment pendimethalin at 1.0 kg ha⁻¹ at 2 DAS + bispyribac sodium at 25 g ha⁻¹ at 20 DAS (T₄).

How to Cite:

Chakraborti, M., Duary, B., Datta, M., 2021. Effect of different weed management practices on the weed population dynamics of direct seeded upland rice under Tripura condition. *Innovative Farming* 6(1), 06-11.

INTRODUCTION

Rice (*Oryza sativa* L.) is the most important staple food for more than half of the world's population, including regions of high population density and rapid growth. India has the largest area among rice growing countries and stands second in production. Expansion in the irrigated area, introduction of early maturing rice cultivars, availability of selective

herbicides for weed management together with increasing transplanting cost and declining profitability of transplanted rice production system have encouraged rice farmers to shift from transplanting to direct seeding (Subbaiah *et al.*, 1999). Globally, actual rice yield losses due to pests have been estimated at 40%, of which weeds have

the highest loss potential (32%). The worldwide estimated loss in rice yield from weeds is around 10% of the total production (Oerke and Dehne, 2004). However, for cultivation of direct-seeded rice, weeds are a major hurdle as nearly all *Kharif* season weeds depending upon seed bank in the field infest this crop.

MATERIALS AND METHODS

A field experiment was conducted at Krishi Vigyan Kendra, South Tripura during the *kharif* (wet) season of 2013 and 2014 to study the influence of different weed management practices on weed population dynamics of direct seeded upland rice under Tripura condition. Twelve treatments, viz., pendimethalin at 1.0 kg ha⁻¹ at 2 DAS (T₁), bispyribac sodium at 25 g ha⁻¹ at 25 DAS (T₂), pendimethalin at 1.0 kg ha⁻¹ at 2 DAS + one hand weeding at 30 DAS (T₃), pendimethalin at 1.0 kg ha⁻¹ at 2 DAS + bispyribac sodium at 25 g ha⁻¹ at 20 DAS (T₄), metsulfuron methyl + chlorimuron ethyl (Almix) at 4 g at 10 DAS followed by bispyribac sodium at 25 g at 20 DAS (T₅), pyrazosulfuron ethyl at 25 g ha⁻¹ at 3 DAS followed by bispyribac sodium at 25 g at 20 DAS (T₆), fenoxaprop-p-ethyl at 60 g ha⁻¹ + metsulfuron methyl + chlorimuron ethyl (Almix) at 4 g ha⁻¹ at 15 DAS (T₇), stale seed bed + smother crop (cowpea) in between two rows of rice (T₈), stale seed bed + one hand weeding at 30 DAS (T₉), sesbania (broadcast) @ 25 kg ha⁻¹ during sowing of rice + 2,4-D at 500 g ha⁻¹ at 25 DAS (T₁₀), hand weeding at 15, 30 and 45 DAS (T₁₁), weedy check (T₁₂) were assigned in a randomized block design replicated thrice. Rice variety NDR-97 was used for the experimental purpose with recommended package of practices.

The upland rice was fertilized as per package of practices recommended (80:40:40 NPK ha⁻¹). Five tonnes of Farm Yard Manure were applied at the time of field preparation for both the crop. Chemical fertilizers were applied to meet 60 kg nitrogen in the form of urea, 40 kg phosphorus in the form of single superphosphate and 40 kg potassium in the form of muriate of potash in the rice.

Weed counts at different stages (15, 30, 60 and at harvest stage) was taken by placing quadrat at random three sites in each plot and calculating the

average. Data on yield attributes, grain yield, straw yield of rice were taken. Economics of different weed management practices was also studied. The data generated from the experiment were subject to analysis of variance (ANOVA) as applied to randomized block design described by Cochran and Cox (1965).

RESULTS AND DISCUSSION

The experimental field was infested with different weed flora namely *Amaranthus viridis*, *Oldenlandia corymbosa*, *Spilanthes acmella*, *Ludwigia parviflora*, *Cleome rutidosperma*, *Malvestrum coromondaliense* among the broad leaf weed, *Digitaria sanguinalis* among grasses and *Cyperus iria* among sedges. Similar weed flora in direct seeded rice was also reported by many researchers like Duary *et al.* (2005), Maity and Mukherjee (2009), Chauhan and Opena (2012), and Kashid *et al.* (2015).

Effect on Weed Density

The effect of various weed management practices on weed density of grassy, broad leaved weed and sedges at different stages showed highly significant during both the years. Among the various treatment tested, all weed management practices resulted in significant reduction in weed density as compared to weedy check. At 30 DAS, the highest number of grassy, broad leaved and sedges were recorded with weedy check treatment. Among the other treatments, pendimethalin at 1.0 kg ha⁻¹ at 2 DAS + bispyribac sodium at 25 g ha⁻¹ at 20 DAS (T₄) recorded the lowest number of all weed in both the years but in the second year this particular treatment was at par with pendimethalin at 1.0 kg ha⁻¹ + one hand weeding at 30 DAS (T₃) (Table 1). There were no remarkable changes in the weed density between two years.

At 60 DAS, hand weeding thrice at 15, 30 and 45 DAS (T₁₁) recorded the lowest population of total weed. Among the other treatments, pendimethalin at 1.0 kg ha⁻¹ + one hand weeding at 30 DAS (T₃) registered the lowest population of total weed and was statistically at par with pendimethalin at 1.0 kg ha⁻¹ at 2 DAS + bispyribac sodium at 25 g ha⁻¹ at 20 DAS (T₄) treatment during both the years. All the

treatments were significantly superior to the weedy check (Table 2).

At 90 DAS, weed density varied significantly with different treatments as evident from the data presented in table 3. The highest population of weeds was recorded with weedy check (T₁₂). The lowest population of weeds was recorded in the

treatment hand weeding at 15, 30 and 45 DAS (T₁₁) in both the years. Among the other treatments, pendimethalin at 1.0 kg ha⁻¹ + one hand weeding at 30 DAS (T₃) registered the lowest number of total weed and was at par with pendimethalin at 1.0 kg ha⁻¹ at 2 DAS + bispyribac sodium at 25 g ha⁻¹ at 20 DAS (T₄) during both the years (Table 3).

Table 1: Effect of treatments on weed density at 30 DAS

Treatment	Weed density (No. m ⁻²) at 60 DAS							
	2013				2014			
	Grasses	BLW	Sedges	Total	Grasses	BLW	Sedges	Total
T ₁	3.29 (10.33)	2.47 (5.66)	3.93 (15.00)	5.61 (31.00)	3.08 (9.00)	2.18 (4.33)	3.71 (13.33)	5.20 (26.66)
T ₂	4.38 (18.66)	3.72 (13.33)	2.90 (8.00)	6.36 (40.00)	4.18 (17.00)	3.53 (12.00)	2.67 (6.66)	6.01 (35.67)
T ₃	3.17 (9.66)	2.48 (5.66)	3.53 (12.00)	5.27 (27.33)	2.02 (3.66)	1.95 (3.33)	3.28 (10.33)	4.22 (17.33)
T ₄	2.47 (5.66)	2.60 (6.33)	3.07 (9.00)	4.62 (21.00)	2.18 (4.33)	2.26 (4.66)	2.84 (7.66)	4.12 (16.66)
T ₅	3.97 (15.33)	2.85 (7.66)	2.47 (5.66)	5.40 (28.66)	3.75 (13.66)	2.73 (7.00)	2.26 (4.66)	5.08 (25.33)
T ₆	3.07 (9.00)	2.96 (8.33)	3.18 (9.66)	5.24 (27.00)	2.79 (7.33)	2.79 (7.33)	2.91 (8.00)	4.80 (22.66)
T ₇	2.72 (7.00)	3.62 (12.66)	2.47 (5.66)	5.08 (25.33)	2.53 (6.00)	3.34 (10.66)	2.11 (4.00)	4.60 (20.66)
T ₈	3.29 (10.33)	3.58 (12.33)	3.33 (10.66)	5.81 (33.33)	2.96 (8.33)	3.33 (10.66)	3.18 (9.66)	5.40 (28.66)
T ₉	3.22 (10.00)	3.33 (10.66)	3.58 (12.33)	5.79 (33.00)	3.01 (8.66)	3.06 (9.00)	3.33 (10.66)	5.37 (28.33)
T ₁₀	4.98 (24.33)	3.44 (11.33)	3.85 (14.33)	7.11 (50.00)	4.77 (22.33)	3.13 (9.33)	3.58 (12.33)	6.67 (44.00)
T ₁₁	0.71 (0)	0.71 (0)	0.71 (0)	0.71 (0)	0.71 (0)	0.71 (0)	0.71 (0)	0.71 (0)
T ₁₂	5.34 (28.00)	5.30 (27.66)	4.52 (20.00)	8.73 (75.66)	5.24 (27.00)	5.18 (26.33)	4.34 (18.33)	8.49 (71.66)
S.Em (±)	0.17	0.14	0.15	0.14	0.18	0.15	0.16	0.15
CD (P=0.05)	0.50	0.42	0.46	0.43	0.53	0.46	0.47	0.44
CV (%)	8.75	8.11	8.68	4.66	10.14	9.67	9.73	5.18

The data on grain yield and straw yield of rice varied significantly among the treatments in both the years (Table 4). There was a drastic reduction in grain yield in the weedy check plot (T₁₂) and it was to the tune of 83.19% in the first year and 79.44% in the second year. Similar results were observed by Naresh *et al.* (2011) and Mathew *et al.* (2013).

Among the various treatment, hand weeding at 15, 30 and 45 DAS (T₁₁) recorded the highest grain and straw yield of upland rice and it was statistically at par with pendimethalin at 1.0 kg ha⁻¹ + one hand weeding at 30 DAS (T₃) and pendimethalin at 1.0 kg ha⁻¹ at 2 DAS + bispyribac sodium at 25 g ha⁻¹ at 20 DAS (T₄) during both the years (Table 4). The



competition between rice and weed for nutrient, water, light and space was less under the above treatments, which facilitated greater utilization of sun light, higher synthesis of carbohydrate and better partitioning of photosynthates towards grain formation and ultimately leading to higher grain

yield of rice. Sardana *et al.* (2004) reported that integrated weed management resulted in lower weed dry matter and higher number of panicles, panicle length and number of grains panicle⁻¹, 1000 grain weight and grain yield. All other treatments were significantly superior than the weedy check (T₁₂).

Table 2: Effect of treatments on weed density at 60 DAS

Treatment	Weed density (No. m ⁻²) at 60 DAS							
	2013				2014			
	Grasses	BLW	Sedges	Total	Grasses	BLW	Sedges	Total
T ₁	16.33	18.00	25.67	60.00	15.00	16.67	24.33	56.00
T ₂	22.33	22.67	12.33	57.33	20.67	21.00	10.67	52.33
T ₃	10.33	9.00	13.00	32.33	8.67	7.67	11.33	27.67
T ₄	10.33	10.00	13.33	33.67	10.00	8.67	12.67	31.33
T ₅	21.00	21.67	12.67	55.33	19.33	20.00	11.00	50.33
T ₆	13.00	17.67	18.33	49.00	11.33	16.00	16.33	43.67
T ₇	18.33	19.67	17.67	55.67	16.33	17.67	15.67	49.67
T ₈	30.33	20.00	28.00	78.33	28.33	18.33	26.33	73.00
T ₉	22.33	21.67	22.00	66.00	20.33	20.33	20.33	61.00
T ₁₀	32.00	17.00	28.00	77.00	29.67	15.33	26.33	71.33
T ₁₁	8.00	6.67	4.67	19.33	7.00	7.00	3.33	17.33
T ₁₂	42.33	34.33	35.67	112.33	40.67	32.33	34.33	107.33
S.Em (±)	1.27	1.11	1.22	1.41	1.21	1.04	1.19	1.41
CD (P=0.05)	3.74	3.26	3.58	4.14	3.57	3.06	3.49	4.14
CV (%)	10.75	10.59	10.98	4.22	11.13	10.81	11.63	4.58

Table 3: Effect of treatments on weed density at 90 DAS

Treatment	Weed density (No. m ⁻²) at 60 DAS							
	2013				2014			
	Grasses	BLW	Sedges	Total	Grasses	BLW	Sedges	Total
T ₁	16.67	17.67	26.67	61.00	15.00	16.33	25.00	56.33
T ₂	22.67	22.67	16.67	62.00	21.33	21.00	15.00	57.33
T ₃	9.67	10.00	12.33	32.00	7.67	9.00	11.33	28.00
T ₄	9.33	11.67	13.00	34.00	9.33	9.67	11.43	30.33
T ₅	20.67	19.33	12.00	52.00	19.33	17.33	10.33	47.00
T ₆	18.33	18.67	20.67	57.67	17.67	17.00	19.00	53.67
T ₇	21.00	20.67	18.00	59.67	19.33	19.00	16.33	54.67
T ₈	32.00	22.67	29.33	84.00	30.33	21.00	27.33	78.67
T ₉	29.00	23.67	23.67	76.33	27.33	22.00	21.67	71.00
T ₁₀	35.33	20.67	29.33	85.33	34.00	19.00	27.33	80.33
T ₁₁	8.33	8.33	4.33	21.00	7.33	7.00	4.00	18.33
T ₁₂	41.67	35.33	37.33	114.33	40.67	34.00	35.33	110.00
S.Em (±)	1.25	1.04	1.05	1.95	1.04	1.05	1.08	1.76
CD (P=0.05)	3.68	3.06	3.09	5.73	3.07	3.09	3.19	5.18
CV (%)	9.85	9.40	9.02	5.49	8.73	10.33	10.09	5.35

The data on net return and return per rupee invested expressed that pendimethalin at 1.0 kg ha⁻¹ at 2 DAS + bispyribac sodium at 25 g ha⁻¹ at 20 DAS (T₄) recorded the highest net return (Rs. 23,847.00 in the first year and Rs. 26,010.00 in the second year) and return rupee⁻¹ invested (2.02 and 2.11) in both the years and was found to be the most remunerative weed management practices (Table 4). The herbicide pendimethalin at 1.0 kg ha⁻¹ + one hand weeding at 30 DAS (T₃) registered net return of Rs. 21,427.00 and Rs. 25,573.00 in the first year and second year

respectively and was the next best treatment. This is simply due to higher grain and straw yield of the crop obtained from these treatments and comparatively lower cost involved in the cultivation of crop under these treatments. Similar result was reported by Yakadri *et al.* (2016). Weedy check (T₁₂) fetched the negative net return. This is due to greater competition between rice and weed which led to poor growth of the crop, lower grain and straw yield. This again emphasized the importance of weed management in profitable rice production.

Table 4: Effect of weed management practices on grain yield, straw yield and harvest index of direct seeded upland rice

Treatment	Grain yield		Straw yield		Economics			
	(t ha ⁻¹)		(t ha ⁻¹)		Net return		Return Rupee ⁻¹ invested	
	2013	2014	2013	2014	2013	2014	2013	2014
T ₁	2.15	2.36	3.89	4.21	11337	14297	1.55	1.70
T ₂	2.21	2.26	3.65	4.22	10650	11830	1.49	1.54
T ₃	3.30	3.59	5.03	5.28	21427	25573	1.81	1.96
T ₄	3.26	3.41	4.87	5.17	23847	26010	2.02	2.11
T ₅	2.49	2.65	3.72	4.65	13488	16542	1.60	1.73
T ₆	2.71	2.80	4.04	4.81	16637	18490	1.73	1.82
T ₇	1.89	1.98	3.82	3.98	7908	9188	1.39	1.45
T ₈	1.86	2.01	3.79	4.01	4493	6613	1.19	1.28
T ₉	1.86	2.10	3.71	3.89	1857	5203	1.07	1.20
T ₁₀	1.81	1.93	3.75	3.74	7490	9083	1.38	1.46
T ₁₁	3.45	3.60	5.16	5.43	17620	19713	1.54	1.61
T ₁₂	0.58	0.74	2.90	2.80	-5347	-3450	0.66	0.78
S.Em (±)	0.11	0.13	0.22	0.17	1609	1783	0.07	0.08
CD (P=0.05)	0.34	0.37	0.64	0.49	4719	5230	0.22	0.23
CV (%)	8.62	8.93	9.40	6.70	25.45	23.29	8.39	8.95

CONCLUSION

In conclusion, the field experiment conducted at Krishi Vigyan Kendra, South Tripura, highlights the crucial role of effective weed management in enhancing the productivity and profitability of direct-seeded upland rice. Among the various treatments evaluated, the combination of pendimethalin at 1.0 kg ha⁻¹ at 2 DAS followed by bispyribac sodium at 25 g ha⁻¹ at 20 DAS emerged as the most effective and economically beneficial strategy. This treatment consistently resulted in the lowest weed density, highest grain yield and superior economic returns. Hand weeding at multiple intervals also proved effective but is less

economically viable due to higher labor costs. The findings underscore the importance of integrated weed management practices, combining chemical and manual methods, to achieve optimal crop performance and profitability in upland rice cultivation under Tripura conditions.

Conflict of Interest

The authors declare no conflict of interest.

REFERENCES

Chauhan, B.S., Opena, J., 2012. Effect of tillage systems and herbicides on weed emergence, weed growth, and grain yield in dry-seeded rice

- systems. *Field Crops Research* 137, 56-69. DOI: <https://doi.org/10.1016/j.fcr.2012.08.016>.
- Cochran, W.G., Cox, G.M., 1965. Experimental Design. John Willey and Sons Inc. New York.
- Duary, B., Hossain, A., Mondal, D.C., 2005. Integrated weed management in direct seeded dry sown rice in lateritic belt of West Bengal. *Indian Journal of Weed Science* 37(1/2), 101-102.
- Kashid, N.V., Barhate, K.K., Bodake, P.S., 2015. Management of weeds in direct seeded rice. *Indian Journal of Weed Science* 47(2), 110-112.
- Maity, S.K., Mukherjee, P.K., 2009. Integrated weed management practices in dry direct-seeded summer rice (*Oryza sativa*). *Indian Journal of Agricultural Sciences* 79(12), 976-979.
- Mathew, R., Raj, S.K., Jose, N., Leenakumary, S., 2013. Comparative efficacy of penoxsulam and pyrazosulfuron ethyl for weed control in direct - seeded puddle rice (*Oryza sativa*). *Indian Journal of Agricultural Sciences* 83(12), 1420-1422.
- Naresh, R.K., Gupta, R.K., Singh, R.V., Singh, D., Singh, B., Prakash, S., Misra, S.P., Rathi, R.C., Bhan, S., 2011. Promotion of integrated weed management for direct-seeded rice in North West India. *Progressive-Agriculture* 11(2), 215-232.
- Oerke, E.C., Dehne, H.W., 2004. Safeguarding production losses in major crops and the role of crop protection. *Crop Protection* 23(4), 275-285. DOI: <https://doi.org/10.1016/j.cropro.2003.10.001>.
- Sardana, V., Mahajan, G., Brar, A.S., Gill, M.S., 2004. Agronomic manipulations for higher productivity of direct seeded rice (*Oryza sativa* L.). *Haryana Journal of Agronomy* 20(1/2), 71-74.
- Subbaiah, S.V., Singh, S.P., Kumar, R.M., Padmaja, K., 1999. Drum seeding technology as a substitute for transplanting in India. In: 2nd IRRE-CREMNET Workshop Cum Group Meeting, 24th-27th August, 1999. Soil and Water Management Research Institute, Thanjavur, Tamil Nadu, India.
- Yakadri, M., Madhavi, M., Ramprakash, T., Leela, R., 2016. Herbicide combinations for control of complex weed flora in transplanted rice. *Indian Journal of Weed Science* 48(2), 155-157.
